**Open Access** 

# The successful treatment of a Gustilo– Anderson type IIIc distal leg injury with a large bone defect in elderly patient with severe osteoporosis: a case report

Koji Nozaka<sup>1\*</sup>, Naohisa Miyakoshi<sup>1</sup>, Motoki Mita<sup>1</sup> and Yoichi Shimada<sup>1</sup>

# Abstract

**Background** Gustilo–Anderson type IIIc tibial open fracture with large bone defects in elderly patients with severe osteoporosis is a rare injury that may be a challenging clinical scenario.

**Case presentation** This study presents the case of a 68-year-old Japanese man who sustained a Gustilo–Anderson type IIIc open tibial fracture with a large bone defect. The patient had severe osteoporosis and the bone was contaminated; therefore, we determined that the bone could not be returned to the tibia. The patient underwent acute limb shortening and gradual lengthening with an Ilizarov external fixator combined with low-intensity pulsed ultrasound and teriparatide administration for limb reconstruction, which allowed immediate full weight-bearing capacity. The fixator was removed at 12 months postoperatively, and by this time, the fracture had completely healed. At the most recent 5-year follow-up after the injury, the patient reported full weight-bearing capacity without walking aids and had full knee and ankle range of motion.

**Conclusions** To the best of our knowledge, this is the first study to report the use of combined Ilizarov technique, low-intensity pulsed ultrasound, and teriparatide for limb reconstruction of Gustilo–Anderson type Illc open tibial fractures with large bone defects in elderly patients with severe osteoporosis.

Keywords Gustilo-Anderson type IIIc open tibial fracture, Severe osteoporosis, Ilizarov external fixator

# Background

The incidence of open tibial fractures in the elderly is increasing due to the growing elderly population. Open tibial fractures with large bone defects are an increasing problem, and their treatment is challenging. Gustilo– Anderson type IIIc open tibial fractures with large bone defects are less common in elderly patients with severe

kk-nozaka@mue.biglobe.ne.jp

osteoporosis than in young patients. However, the literature on the outcomes of open tibial fractures with large bone defects in elderly patients treated with modern techniques is limited.

# **Case presentation**

A 68-year-old Japanese man was injured while electrically reeling a wire rope on a fishing boat. Shortly after, the patient was transported to our hospital's emergency room, where he was treated for hypertension. The patient had no relevant family or psychosocial history. On physical examination, two large soft-tissue defects were observed in the medial left ankle (Fig. 1). The patient's left foot was pale, and we did not palpate any arteries in



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/. The Creative Commons Public Domain Dedication waiver (http://creativecommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

<sup>\*</sup>Correspondence:

Koji Nozaka

<sup>&</sup>lt;sup>1</sup> Department of Orthopedic Surgery, Akita University Graduate School of Medicine, 1-1-1 Hondo, Akita 010-8543, Japan



Fig. 1 Two large soft-tissue defects in the medial left ankle

his lower legs because of severe ankle pain. Next, pulses from the peroneal, tibialis anterior, and tibialis posterior arteries, as well as the capillary refill, color, and temperature, were compared between the injured and uninjured feet. Then, the paramedics brought in a 75-mm tibia that had fallen out of the body on the fishing boat. The bone was contaminated; therefore, we determined that the bone could not be returned to the tibia (Fig. 2a-c). Three-dimensional (3D) computed tomography (CT) angiography revealed injury to the peroneal, tibialis anterior, and tibialis posterior arteries (Fig. 3a, b). The preoperative Mangled Extremity Severity Score (MESS) was 7 points. We considered limb salvage feasible and performed reconstructive surgery. A total of 6 hours after the injury, we debrided the soft tissues of the injury in the operating room (day 0). After debridement, the softtissue defect measured 40 mm × 80 mm. We decided that acute limb shortening and gradual lengthening would be a reasonable method for rapid revascularization. Accordingly, we shortened the fibula with a 75-mm osteotomy to match the length of the tibia. We resected and prepared a spike in the distal tibial fragment (Fig. 4a). We crimped the fracture sufficiently and fixed it with an Ilizarov external fixator for temporary ankle joint-bridging fixation



**Fig. 3** a 3D computed tomography angiography image with the bone; b 3D computed tomography angiography image without the bone *3D* three-dimensional

(Fig. 4b, c). We reconstructed the anterior and posterior tibial arteries and veins with end-to-end anastomosis (day 0). No severe nerve injury to the lower leg was obs erved.

Revascularization was completed 6 hours after the injury, after which circulation in the injured foot improved. The size of the soft-tissue defect was now  $20 \times 80$  mm. Negative pressure wound therapy was used to address the soft-tissue defect (Fig. 4d). After the patient gained consciousness following anesthesia, ankle joint circulation was good and automatic movement of the ankle joint and toes was possible (Fig. 4e). The patient had normal motor and sensory nerves in his lower leg.



Fig. 2 a The contaminated large bone that fell out of the body; b and c X-ray images taken in the emergency room



**Fig. 4** a Shortened fibula with a 75-mm osteotomy to match the length of the tibia; **b** and **c** the fibula was fixed with an Ilizarov external fixator for temporary ankle joint-bridging fixation; **d** and **e** clinical photograph of the fibula, which was fixed with an Ilizarov external fixator for temporary ankle joint-bridging fixation

Although the patient had mild renal dysfunction, he did not undergo hemofiltration and recovered without permanent organ damage.

A total of 2 days after the injury (day 2), we cleaned the open fracture again, performed additional debridement, and reapplied negative pressure wound therapy. A total of 14 days after the injury (day 14), osteotomy was performed at the proximal tibia with an Ilizarov external fixator for gradual limb lengthening (Fig. 5a). A total of 3 weeks after the injury (day 21), gradual lengthening of the proximal tibia was initiated. We noted that the bone strength was weak at the time of wire insertion; therefore, we measured the bone density and identified primary osteoporosis. A baseline dual-energy X-ray absorptiometry scan showed that the bone mineral density at the femoral neck was 0.441 g/cm<sup>2</sup> and T-Score was -3.3 standard deviation. He was started on a onceweekly injection of 56.5 µg of teriparatide. Furthermore, after osteotomy, treatment with a low-intensity pulsed ultrasound stimulation (LIPUS) device (SAFHS 2000, Exogen, Inc., Piscataway, NJ) was started for 20 minutes/ day at the fracture and osteotomy sites. This device had a frequency of 1.5 MHz, a signal burst width of 200 µs, a signal repetition frequency of 1 kHz, and an intensity of 30 mW/cm<sup>2</sup>. The patient was allowed to walk with full weight-bearing capacity immediately after the surgery (Fig. 5b) (Day 22). He also performed knee and ankle range of motion exercises (day 22). We performed



Fig. 5 a Osteotomy at the proximal tibia with an Ilizarov external fixator for gradual lengthening; b bone lengthening at a rate of 0.75 (0.25 × 3) mm/day; c walking with full weight-bearing capacity immediately after surgery

bone lengthening at a rate of 0.75 ( $0.25 \times 3$ ) mm/day. The patient continued to undergo bone lengthening at home after discharge. The patient visited the outpatient clinic once a month for radiography. The patient was positive about returning to his original job, and was consistently mentally stable. Radiography performed 1 year later revealed good callus formation (Fig. 5c). The fixator was removed after one year. The patient returned to his original job as a fisherman 2 months after the fixator removal. The patient's 14 months of treatment was covered by industrial accident compensation insurance.

At the 5-year follow-up after injury, radiographs showed good callus formation and bone union (Fig. 6a, b, c, and d). The patient was independently mobile, with a knee range of motion of  $0-140^{\circ}$ , ankle dorsiflexion of 5°, and plantar flexion of 50° (Fig. 7a, b, and c). During the 12 months in which the fixator was inserted, there were a few superficial pin-tract infections, which were treated with empirical oral antibiotics and daily pin-tract dressings. A total of 5 years after surgery, the patient returned to work without any problems, with an American Orthopaedic Foot and Ankle Society (AOFAS) score of 95 points. The clinical results, according to the AOFAS score, were excellent in this patient.

## **Discussion and conclusions**

Treatment of high-energy open tibial fractures is challenging. Gustilo-Anderson type IIIc open tibial fractures were mainly treated with amputation in the past [1]. The complication rates are high for Gustilo-Anderson type IIIc open tibial fractures, which are associated with more severe soft-tissue loss and/or arterial injury requiring repair [2]; these complications include secondary amputation, nonunion, infection, and malunion [3-5]. The acceptance rate of amputation is poor in eastern-culture patients, especially in Japanese elderly patients [6]. Adaptation to a prosthesis is relatively easier in young amputees than in elderly amputees [7]. In our patient, there was a bone defect, soft-tissue defect, and peroneal, tibialis anterior, and tibialis posterior artery tears. The advantage of acute shortening is that it is easier to reconstruct the arteries and veins with end-to-end anastomosis.



Fig. 6 a-d X-ray images taken at postoperative 5-year follow-up

Furthermore, soft-tissue defects were smaller in genaral; therefore, function reconstruction was possible in our elderly patient, despite his presentation. In Gustilo-Anderson type IIIb open tibial fractures treated with acute shortening, gradual lengthening is usually subjected to a second-stage procedure performed after union of the fracture [8]. The main disadvantage of this approach is prolonged treatment duration. In Gustilo-Anderson type IIIc open tibial fractures, lengthening is delayed to prevent traction on vascular anastomoses as they heal, which may potentially risk the revascularization procedure. In our study, we performed early lengthening without causing any harm to vascular repair, as lengthening was performed at the proximal tibia most distant from the vascular injury. Management of open injuries of the limbs are challenging, as there are still many gray areas in decision making regarding salvage, timing, and reconstruction type. As a result, there is still an unacceptable rate of secondary amputations, which leads to tremendous waste of resources and psychological devastation of the patient and their family [9]. In addition, limb salvage is more cost effective than amputation and prosthesis use [10, 11]. Our patient achieved a satisfactory functional status and avoided psychological trauma due to amputation; he was able to return to his original position as a fisherman.

It is difficult to reconstruct large bone defects in elderly patients with severe osteoporosis due to decreased bone formation. LIPUS has been used to treat leg lengthening [12, 13]. Intermittent administration of human parathyroid hormone (PTH) has an anabolic effect on the bone in humans and is expected to be a potent agent for fracture healing [14]. Several recent studies have revealed that intermittent treatment with PTH stimulates osteogenesis in experimental fracture healing of cortical bones and that the effects of PTH on cortical bone repair are site-specific. Aspenberg et al., in a prospective randomized double-blind study of conservative fracture treatment for 102 postmenopausal women with distal radial fractures, showed that the time to healing was shorter in patients who received 20 mg teriparatide than in the placebo group patients [15]. Warden et al. reported that teriparatide and LIPUS have contrasting additive rather than synergistic effects during fracture healing [16]. Teriparatide primarily increased the callus bone mineral content without influencing its size, whereas LIPUS increased callus size without influencing the callus bone mineral







Fig. 7 a-c Clinical photograph at postoperative 5-year follow-up

content in rat models [16]. We have reported combined effect of teriparatide and low-intensity pulsed ultrasound for patients with nonunion [17]. Furthermore, we have reported intractable fractures such as pathological fractures in patients with Alagille syndrome or nonunion after ankle fracture for Charcot arthropathy that was treated with LIPUS and an Ilizarov external fixator [18, 19]. Early ambulation and immediate weight-bearing capacity may improve limb circulation and enhance the healing process, based on the fact that the speed of fracture healing is usually proportional to the amount of available circulation to and between fragments [20, 21].

One of the most important advantages of using Ilizarov external fixators is the excellent recorded knee and ankle range of motion within a short time after surgery. Active and passive movements of both the joints were allowed and encouraged during the entire course of treatment immediately after application of the frame. The main disadvantages of Ilizarov external fixators are that they are technically demanding and there is absolute necessity of adequate care of the frame. From our point of view, the fact that the patient could achieve immediate weightbearing capacity and could be discharged and returned to work is adequate justification for this procedure.

#### Abbreviations

LIPUS Low-intensity pulsed ultrasound stimulation PTH Parathyroid hormone

#### Acknowledgements

Not applicable

#### Author contributions

KN performed the surgery. MN helped with surgery and helped to draft the manuscript. MM and YS helped draft the manuscript. All authors read and approved the final manuscript.

#### Funding

The author(s) received no financial support for the research, authorship, and/ or publication of this article.

#### Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

## Declarations

#### Ethics approval and consent to participate

The patient received prior information before providing his written, informed consent in accordance with the Declaration of Helsinki. The article was approved by the ethics committee of Akita University Hospital (registration number 1970).

#### **Consent for publication**

Written informed consent was obtained from the patient for publication of this case report and any accompanying images. A copy of the written consent is available for review by the Editor-in-Chief of this journal.

#### **Competing interests**

All authors declare that they have no competing interests.

Received: 15 September 2021 Accepted: 6 January 2023 Published online: 13 October 2023

#### References

- 1. Quirke TE, Sharma PK, Boss WK Jr, Oppenheim WC, Rauscher GE. Are type IIIC lower extremity injuries an indication for primary amputation. J Trauma. 1996;40(6):992–6. https://doi.org/10.1097/00005373-19960 6000-00023.
- Gustilo RB, Mendoza RM, Williams DN. Problems in the management of type III (severe) open fractures: a new classification of type III open fractures. J Trauma. 1984;24(8):742–6. https://doi.org/10.1097/00005373-198408000-00009.
- Allen MJ, Nash JR. loannidies TT, Bell PR: major vascular injuries associated with orthopaedic injuries to the lower limb. Ann R Coll Surg Engl. 1984;66(2):101–4.
- Howe HR Jr, Poole GV Jr, Hansen KJ, Clark T, Plonk GW, Koman LA, Pennell TC. Salvage of lower extremities following combined orthopedic and vascular trauma. A predictive salvage index. Am Surg. 1987;53(4):205–8.
- DiChristina DG, Riemer BL, Butterfield SL, Burke CJ 3rd, Herron MK, Phillips DJ. Femur fractures with femoral or popliteal artery injuries in blunt trauma. J Orthop Trauma. 1994;8(6):494–503.
- Tunali O, Saglam Y, Balci HI, Kochai A, Sahbaz NA, Sayin OA, Yazicioglu O. Gustilo type IIIC open tibia fractures with vascular repair: minimum 2-year follow-up. Eur J Trauma Emerg Surg. 2017;43(4):505–12. https://doi.org/ 10.1007/s00068-016-0689-y.
- Parmaksizoglu F, Koprulu AS, Unal MB, Cansu E. Early or delayed limb lengthening after acute shortening in the treatment of traumatic belowknee amputations and Gustilo and Anderson type IIIC open tibial fractures: the results of a case series. J Bone Joint Surg Br. 2010;92(11):1563–7. https://doi.org/10.1302/0301-620X.92B11.23500.
- Yokoyama K, Itoman M, Nakamura K, Uchino M, Tsukamoto T, Suzuki T. Primary shortening with secondary limb lengthening for Gustilo IIIB open tibial fractures: a report of six cases. J Trauma. 2006;61(1):172–80. https:// doi.org/10.1097/01.ta.0000225049.41381.9d.
- Rajasekaran S, Sabapathy SR, Dheenadhayalan J, Sundararajan SR, Venkatramani H, Devendra A, Ramesh P, Srikanth KP. Ganga hospital open injury score in management of open injuries. Eur J Trauma Emerg Surg. 2015;41(1):3–15. https://doi.org/10.1007/s00068-014-0465-9.
- Hertel R, Strebel N, Ganz R. Amputation versus reconstruction in traumatic defects of the leg: outcome and costs. J Orthop Trauma. 1996;10(4):223–9. https://doi.org/10.1097/00005131-199605000-00001.
- Williams MO. Long-term cost comparison of major limb salvage using the Ilizarov method versus amputation. Clin Orthop Relat Res. 1994;301:156–8.
- Gebauer D, Correll J. Pulsed low-intensity ultrasound: a new salvage procedure for delayed unions and nonunions after leg lengthening in children. J Pediatr Orthop. 2005;25(6):750–4. https://doi.org/10.1097/ 01bpo.0000173245.12184.7e.
- Sato W, Matsushita T, Nakamura K. Acceleration of increase in bone mineral content by low-intensity ultrasound energy in leg lengthening. J Ultrasound Med. 1999;18(10):699–702. https://doi.org/10.7863/jum.1999. 18.10.699.
- Nozaka K, Miyakoshi N, Kasukawa Y, Maekawa S, Noguchi H, Shimada Y. Intermittent administration of human parathyroid hormone enhances bone formation union at the site of cancellous bone osteotomy in normal and ovariectomized rats. Bone. 2008;42(1):90–7. https://doi.org/ 10.1016/j.bone.2007.08.041.
- Aspenberg P, Genant HK, Johansson T, Nino AJ, See K, Krohn K, García-Hernández PA, Recknor CP, Einhorn TA, Dalsky GP, Mitlak BH, Fierlinger A, Lakshmanan MC. Teriparatide for acceleration of fracture repair

in humans: a prospective, randomized, double-blind study of 102 postmenopausal women with distal radial fractures. J Bone Miner Res. 2010;25(2):404–14. https://doi.org/10.1359/jbmr.090731.

- Warden SJ, Komatsu DE, Rydberg J, Bond JL, Hassett SM. Recombinant human parathyroid hormone (PTH 1–34) and low-intensity pulsed ultrasound have contrasting additive effects during fracture healing. Bone. 2009;44(3):485–94. https://doi.org/10.1016/j.bone.2008.11.007.
- Nozaka K, Shimada Y, Miyakoshi N, Yamada S, Hongo M, Kasukawa Y, Saito H, Kijima H. Combined effect of teriparatide and low-intensity pulsed ultrasound for nonunion: a case report. BMC Res Notes. 2014;27(7):317. https://doi.org/10.1186/1756-0500-7-317.
- Nozaka K, Shimada Y, Miyakoshi N, Yamada S, Kasukawa Y, Noguchi A. Pathological fracture of the femur in Alagille syndrome that was treated with low-intensity pulsed ultrasound stimulation and an Ilizarov ring fixator: a case report. BMC Musculoskelet Disord. 2014;8(15):225. https://doi. org/10.1186/1471-2474-15-225.
- Nozaka K, Shimada Y, Kimura Y, Yamada S, Kashiwagura T, Sakuraba T, Wakabayashi I. Successful treatment of nonunion with an Ilizarov ring fixator after ankle fracture for Charcot arthropathy: a case report. BMC Res Notes. 2014;7(7):503. https://doi.org/10.1186/1756-0500-7-503.PMID:25103697Fr eePMCarticle.
- El-Sayed M, Atef A. Management of simple (types A and B) closed tibial shaft fractures using percutaneous lag-screw fixation and Ilizarov external fixation in adults. Int Orthop. 2012;36(10):2133–8. https://doi.org/10. 1007/s00264-012-1620-8.
- Nozaka K, Miyakoshi N, Saito H, Chida S, Tsuchie H, Shimada Y. Effectiveness of Ilizarov external fixation in elderly patients with pilon fractures. J Orthop Sci. 2021;26(2):254–60. https://doi.org/10.1016/j.jos.2020.02.018.

## **Publisher's Note**

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

#### Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

#### At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

